

Secukinumab effect in two types of arthritis seen in children and adolescents at two years

Full abstract title: Efficacy and Safety of Secukinumab in Enthesitis-related Arthritis and Juvenile Psoriatic Arthritis: Primary Results from a Randomised, Double-blind, Placebo-controlled, Treatment Withdrawal, Phase 3 Study (JUNIPERA)

Authors: Ruperto N, Foeldvari I, Alexeeva E, et al. Date: June 2021

Please note that this summary only contains information from the full EULAR 2021 scientific abstract and selected supporting references. The results of this study may not reflect those of other studies. This summary is not intended to provide medical advice.


To investigate if secukinumab is an effective and safe treatment for juvenile psoriatic arthritis

Why was this study done?

JIA is the most common type of arthritis in children and teenagers, beginning before the age of 16.1 Symptoms of JIA can begin from as young as two years1 and gradually worsen over time. All types of JIA

(JPsA) and enthesitis-related arthritis (ERA), two subtypes of juvenile idiopathic arthritis (JIA).

can dramatically impact a child's life, causing pain, tenderness and stiffness in joints all over the body.² There are seven subtypes of JIA*. This study focused on two types: JPsA and ERA. JIA



JIA is an autoimmune disease,² meaning that the body's immune system is overactive, incorrectly attacking normal cells and causing unwanted inflammation. Inflammation can be good (eg, in fighting infection), but it can also become a problem if it occurs

smoothly against each other) is gradually destroyed.2 Idiopathic means that the disease has no known cause. It is thought that certain genes inherited from your parents may be activated by a virus, bacteria or other factors. There is no evidence that external factors such as allergies, toxins in the

In JIA, the overactive immune system causes joints to become swollen and painful, as inflammation kicks in, and cartilage (the tissue between joints that helps bones slide

JIA is the most common childhood rheumatic disease,¹ affecting approximately 2 million children worldwide.³ The subtypes of JPsA and ERA each account for up to 11% of total JIA cases.1



Secukinumab is a type of medication called a biologic. It works to reduce inflammation by blocking one of the molecules that causes inflammation to occur.4 Therefore, it could help relieve the symptoms of JIA.



JPsA and ERA.

secukinumab treatment.

What did this study look at?

The study looked at the effect of secukinumab on the symptoms of two subtypes of JIA:

To observe the effect secukinumab had on joint symptoms, the time it took for joint symptoms to worsen (known as time to flare⁵) was measured between three months (12 weeks) and two years (104 weeks) of


50%

more than needed or without a good reason.

environment or food can cause the disease.2

To check if any improvement was because of secukinumab, results were compared with the time to flare in patients when given a 'dummy' injection containing no active ingredient, called a placebo.

100% 90% The study also measured how much secukinumab improved the severity of joint 70%



In treatment period 1.

response at the end of

this period progressed

104

symptoms. To do this, a specific method

improvement seen in joint pain, mobility and

stiffness. This was referred to as JIA ACR 30/50/70/90/100, meaning a 30, 50, 70, 90


was used to calculate the amount of

or 100% improvement in symptoms.

all patients received secukinumab for 12 weeks. Patients achieving at JPsA and ERA least JIA ACR 30

patients, aged

2 to 17 years



In treatment period 2,

patients were allocated to one

of two arms: secukinumab

and response was observed

75 mg/150 mg or placebo

through Week 104

those given placebo, or a 'dummy' treatment containing no active ingredient, with a 72% reduction in risk of flare

continued to be treated with secukinumab to two years, it was shown that it took significantly longer for their joint symptoms to get worse (time to flare) compared with

JIA ACR 30 response

JIA ACR 70 response

placebo at two years.

9 in 10 children in the secukinumab group had achieved a 30% improvement in joint symptoms with secukinumab treatment compared with 6 in 10 children in the placebo group

Number of flares in treatment period 2

8 in 10 children in the secukinumab group had

achieved a 50% improvement in joint symptoms

with secukinumab treatment compared with

6 in 10 children in the placebo group

21

JIA ACR 50 response

10

flares

More patients achieved and maintained JIA ACR 30 and JIA ACR 70 with secukinumab compared to

The study showed that secukinumab significantly reduces the risk of joint symptoms worsening in

Why does this matter?

children with JPsA and ERA for at least two years. Current treatment options for JPsA and ERA are limited and only a minority of patients are able to control their disease. 6 Secukinumab could potentially offer a much-needed treatment.

7 in 10 children in the secukinumab group had achieved a 70% improvement in joint symptoms with

secukinumab treatment compared with 4 in 10 children in the placebo group

Secukinumab was well-tolerated with no new or unexpected side effects.

Glossary

Biologic medicine: a treatment made using living organisms, rather

Arthritis

[arth-rye-tiss]:

stiffness of the joints.

Safety

than being chemically synthesized. **Enthesitis** [en-thee-sye-tiss]:

inflammation of the entheses, the sites where tendons or ligaments insert into the bone.

a disease causing painful inflammation and

JIA ACR 30/50/70/90/100:

[id-ee-oh-path-ic]: of unknown cause.

a way to measure improvement in JIA joint disease symptoms, where each number refers to 30%, 50%, 70%, 90% and 100% improvement from a baseline value, respectively.

Placebo: a substance with no active component which has no therapeutic effect.

Significant(ly):

age of 16.

Juvenile

[joo-vuh-nile]:

statistically, the difference between the groups is unlikely to have occurred by chance. This difference is therefore likely to be related to the medication given to the patients.

for or relating to young people.

Juvenile idiopathic arthritis:

an umbrella term describing seven different types

of arthritis in children and teenagers under the

Who sponsored this study?

sponsored both this study and the writing of this plain language media summary.

arthritis. [Last accessed: June 2021].

Novartis Pharma AG, Basel, Switzerland

Further information

More on this study can be found here: https://clinicaltrials.gov/ct2/show/NCT03031782

References 1. Momah T, Ray L. J Fam Pract. 2019;68: (2) E8-E13.

2. Arthritis Foundation. Juvenile Idiopathic Arthritis (JIA). Available from: https://www.arthritis.org/diseases/juvenile-idiopathic-

3. Dave M, et al. Pediatr Rheumatol Online J. 2020;18:49 4. Blair HA. Drugs. 2019;79:(4)433-443. 5. NHS. A Flare-up of Juvenile Idiopathic Arthritis. Information for young people, parents and carers. Available from: http:// flipbooks.leedsth.nhs.uk/LN000935.pdf. [Last accessed: June 2021].

Glerup M, et al. Arthritis Care Res (Hoboken). 2020;72:(4)507-516.